

**2005 NATIONAL CPM DATA COLLECTION – NATIONAL AND NETWORK PROFILES
for Adult (aged ≥ 18 years) In-Center Hemodialysis Patients**

Network	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	US
# in sample	452	474	479	465	470	477	468	476	475	469	463	446	470	486	479	470	476	484	8479

DIALYSIS ADEQUACY

% Pts with mean spKt/V ≥ 1.2	94	86	91	92	89	91	90	93	94	91	88	93	90	94	91	92	88	88	91
Median spKt/V	1.55	1.50	1.52	1.56	1.51	1.54	1.53	1.53	1.55	1.59	1.52	1.55	1.53	1.62	1.59	1.62	1.52	1.54	1.55
% Pts with mean URR ≥ 65%	91	82	88	89	85	88	88	86	92	89	85	91	86	92	87	89	85	86	88
Median URR %	74	72	72	73	72	73	73	72	73	74	73	73	72	74	74	75	72	73	73
Median blood pump flow (mL/min)	400	400	400	400	400	400	400	402	400	400	404	401	400	400	400	400	400	400	400
Median dialysis session length (min)	210	210	211	227	210	214	210	225	225	220	210	212	220	236	212	233	198	210	213

VASCULAR ACCESS

% Prevalent pts with AVF	49	50	34	38	31	37	43	30	36	36	37	36	35	35	46	57	47	38	39
% Incident pts with AVF	42	40	43	36	24	38	33	32	26	30	31	38	38	45	34	77	61	48	38
% Prevalent pts with AVG	24	25	32	31	39	41	27	44	31	30	32	35	35	44	29	20	33	39	34
% pts with AVG and stenosis monitoring	50	51	57	66	62	71	63	78	77	65	62	77	55	68	61	77	83	69	67
% Prevalent pts with catheter	28	26	34	31	31	22	30	26	33	33	32	29	31	21	26	23	20	23	27
% Prevalent pts with catheter ≥ 90 days	22	19	31	24	23	17	23	18	25	21	25	21	21	16	24	15	15	17	21

2005 NATIONAL CPM DATA COLLECTION – NATIONAL AND NETWORK PROFILES (cont.)
for Adult (aged ≥ 18 years) In-Center Hemodialysis Patients

Network	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	US
ANEMIA MANAGEMENT																			
Median hgb (g/dL)	11.8	12.0	12.0	12.0	12.2	12.1	12.0	12.0	12.1	12.1	12.0	12.0	12.1	12.0	12.2	11.9	12.0	12.1	12.0
% Pts with mean hgb ≥ 11g/dL	80	79	84	85	84	84	84	85	84	83	83	83	80	84	84	82	82	87	83
% Pts with mean hgb 11-12.0 g/dL [^]	38	30	34	41	30	31	36	38	28	32	32	37	30	35	31	38	34	37	34
% Pts with mean hgb < 10 g/dL	7	9	7	5	8	4	6	7	4	8	6	5	7	3	6	3	6	4	6
Median wkly IV epoetin dose units/kg/wk	215	244	231	193	204	203	226	223	187	193	181	195	199	188	164	178	182	187	198
Median wkly SC epoetin dose units/kg/wk	147	175	159	*	*	150	338	*	184	136	183	114	197	98	103	117	142	234	150.0
% Pts rx'd [^] SC epoetin	2	3	11	*	*	3	2	*	3	5	3	5	3	11	6	8	9	11	5

IRON MANAGEMENT

% Pts with mean TSAT ≥ 20%	75	77	74	75	83	83	81	77	78	76	79	74	78	84	81	71	73	85	79
Median TSAT %	24	26	25	25	27	27	28	25	26	26	27	25	26	27	27	25	25	29	26
% Pts with mean ferritin ≥ 100 ng/mL	91	93	95	92	93	95	93	95	97	95	93	95	96	95	93	94	93	93	94
Median ferritin ng/mL	502	502	547	512	524	521	533	530	552	552	510	474	554	578	464	414	467	562	521
% Pts rx'd IV iron	70	65	75	76	67	72	70	71	74	73	69	73	72	69	70	70	71	63	70

[^] Among those patients prescribed epoetin * Value suppressed because n ≤ 10 Note: To convert hemoglobin conventional units of g/dL to SI units (g/L), multiply by 10.

**2005 NATIONAL CPM DATA COLLECTION – NATIONAL AND NETWORK PROFILES (cont.)
for Adult (aged ≥ 18 years) In-Center Hemodialysis Patients**

Network	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	US
ALBUMIN																			
% Pts with mean serum albumin ≥ 4.0/3.7g/dL (BCG/BCP) ^{^^}	30	31	31	31	37	40	33	37	35	36	34	31	38	38	35	35	41	39	36
% Pts with mean serum albumin ≥ 3.5/3.2g/dL (BCG/BCP)	79	77	79	79	83	82	85	81	81	82	80	82	83	83	84	81	82	85	81
Median serum BCG albumin (g/dL)	3.8	3.8	3.8	3.8	3.9	3.9	3.8	3.9	3.8	3.9	3.8	3.8	3.8	3.8	3.9	3.8	3.9	3.9	3.8
Median serum BCP albumin (g/dL)	3.3	3.5	3.8	3.3	*	3.5	3.6	*	3.6	3.6	3.7	3.5	3.8	3.8	3.6	3.4	3.6	3.8	3.6

^{^^}BCG/BCP-Bromcresol Green/Bromcresol Purple Laboratory Methods

* Value suppressed because n ≤ 10

Note: To convert serum albumin conventional units of g/dL to SI units (g/L), multiply by 10.